



12-2020

## **Impact of an Acute Heat Shock During In Vitro Maturation on Interleukin 6 and its Associated Receptor Component Transcripts In Bovine Cumulus-Oocyte Complexes**

Julia Rowinski

*University of Tennessee, [jrowinsk@vols.utk.edu](mailto:jrowinsk@vols.utk.edu)*

Follow this and additional works at: [https://trace.tennessee.edu/utk\\_gradthes](https://trace.tennessee.edu/utk_gradthes)

---

### **Recommended Citation**

Rowinski, Julia, "Impact of an Acute Heat Shock During In Vitro Maturation on Interleukin 6 and its Associated Receptor Component Transcripts In Bovine Cumulus-Oocyte Complexes. " Master's Thesis, University of Tennessee, 2020.

[https://trace.tennessee.edu/utk\\_gradthes/6098](https://trace.tennessee.edu/utk_gradthes/6098)

This Thesis is brought to you for free and open access by the Graduate School at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Masters Theses by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

To the Graduate Council:

I am submitting herewith a thesis written by Julia Rowinski entitled "Impact of an Acute Heat Shock During In Vitro Maturation on Interleukin 6 and its Associated Receptor Component Transcripts In Bovine Cumulus-Oocyte Complexes." I have examined the final electronic copy of this thesis for form and content and recommend that it be accepted in partial fulfillment of the requirements for the degree of Master of Science, with a major in Animal Science.

J. Lannett Edwards, Major Professor

We have read this thesis and recommend its acceptance:

F. Neal Schrick, Kyle McLean

Accepted for the Council:

Dixie L. Thompson

Vice Provost and Dean of the Graduate School

(Original signatures are on file with official student records.)

**Impact of an Acute Heat Shock During In Vitro Maturation on  
Interleukin 6 and its Associated Receptor Component Transcripts In  
Bovine Cumulus-Oocyte Complexes**

A Thesis Presented for the  
Master of Science  
Degree  
The University of Tennessee, Knoxville

Julia Rose Rowinski

December 2020

## ACKNOWLEDGMENTS

I would like to first extend many thanks to Dr. Edwards for mentoring me over the last 2 years. Her attention to detail will be a skill that I can utilize not only in my career but life as well. I appreciate that she truly cares about her students and always watches out for them. The opportunities given to me in the last two years not only in the lab but on farm will remain with me for a lifetime.

Along with Dr. Edwards, I wanted to thank my committee members Dr. Schrick and Dr. McLean. I know time is precious and I appreciate the time they spent with me in meetings and answering questions throughout my time at UT. Towards that end, I would like to extend a massive thank you to Louisa and Becca for all their help in the lab. Thank you to Louisa for not only taking the time to show me the various skill set I would need for this project and its analyses but for encouraging me to grow as a scientist in every way. I appreciate that she always made sure I got home okay when my shift of the project sampling were done. Thank you to Becca for teaching me the ins and outs of IVP and bestowing her “queen of the lab” knowledge on me, even if that means I will forever take bread out of the bag in a sterile manner. I appreciate that she took the time and saw what I could be by never accepting a mediocre job from me. I am so thankful to have had the opportunity to work alongside such intelligent strong scientists.

Thank you to Dr. Schneider for teaching me the skills I needed to run SAS not only in a classroom but outside of as well. I appreciate showing me the countless ways to analyze to make sure the most appropriate was used and for referring me to Xiaocun Sun. Which leads me to Xiaocun Sun from Research Computing Support with OIT who showed me how to analyze output as well as export tables and graphs from JMP. I appreciate that she was always quick to respond to emails and willing to set up a zoom meeting when it still did not make sense.

Thank you to my fellow lab mates I could not have made it through without you. Hunter Liles I am so glad that you were my graduate school sister and we made it through together. We were always able to count on one another to be there to bounce ideas off of as well as celebrate wins and work through losses. Thank you for bringing me coffee during collections and thank you to Mark Phillips for running a sandwich and chips to me so I could make it through the final hours of the study. Thank you to Dr. Jessica Klabnik for being the older sister I have never had and keeping both Hunter and I in line and motivated to finish. I hope she will remember that I will always be there to support her as she finishes up. I truly and honestly do not think I would have made it through without either of them.

As always this would not have been possible without the love and support of my parents Christina and Stephen Rowinski. Even if they did not always understand exactly what I was doing they continued their support and always knew I could do it. Thank you to Auntie Roxy for her inspiration to study reproductive physiology and even if embryologist does not end up as my title I am forever grateful for the inspiration to be able to do whatever it is my heart desires.

I would like to also take time to thank Evan Prosch, while he may have come in late to the game his constant support for me was vital to my completion of this degree program. I appreciate dinners he made and early morning calls to make sure I was up. He was and is a constant reminder that I can do this and tomorrow is a new day.

Finally, a sincere thank you to the department of animal science of University of Tennessee Institute of Agriculture for the financial support as I completed my degree program. This project was supported by Agriculture and Food Research Initiative Competitive Grant no. 2016-67015-24899 from the USDA National Institute of Food and Agriculture, the state of Tennessee through UT AgResearch, East Tennessee Research and Education Center, the Department of Animal Science, and the USDA National Institute of Food and Agriculture, Hatch Project No. 1015707 and 1022068.

## ABSTRACT

An acute heat stress event after LH surge increased intrafollicular interleukin 6 (IL6) levels in cows that became hyperthermic. To examine direct consequences of the physiologically-relevant elevated temperature of 41.0°C on the cumulus-oocyte complex (COC), *IL6* transcript abundance and related receptor components were evaluated throughout *in vitro* maturation. Because IL6 and LIF share a signal transducer, *LIF* was also examined. Abundance of *IL6* was interactively influenced by temperature and time ( $P < 0.0001$ ). Heat-induced increases in *IL6* were first noted at 4 hIVM; peak levels occurred at 4.67 versus 6.44 hIVM for 41.0 and 38.5°C COCs, respectively (SEM = 0.23;  $P < 0.001$ ). Abundance of *IL6R* ( $P = 0.05$ ) and *IL6ST* ( $P = 0.002$ ) were also interactively influenced by temperature and hIVM. Peak *IL6ST* levels occurred at 6.95 versus 8.29 hIVM for 41.0 and 38.5°C, respectively (SEM = 0.23;  $P < 0.01$ ). Transcript for *LIF* differed over time ( $P < 0.0001$ ) but was not affected by 41.0°C exposure. Blastocyst development after performing IVF was not affected by 41.0°C exposure for 4 or 6 h. Progesterone released into maturation medium by COCs was interactively influenced by temperature and hIVM ( $P = 0.0116$ ). When limiting analysis to when *IL6* was temporally produced, progesterone levels were only impacted by time and temperature. Heat-induced shift in the temporal production of *IL6* along with its impact on progesterone likely cooperate in heat-induced hastening of meiotic progression described by others.

### **Keywords:**

Cumulus-oocyte complex, heat stress, Interleukin 6, oocyte maturation, progesterone

## TABLE OF CONTENTS

Chapter One Introduction and Review of Pertinent Literature.....	6
Chapter Two Materials and Methods .....	8
Collection and <i>in vitro</i> maturation of cumulus-oocyte complexes.....	8
Study one: Interleukin 6, IL6 Receptor, Signal Transducer and LIF transcripts during <i>in vitro</i> maturation in COCs matured at 38.5 or 41.0°C .....	8
Total RNA isolation, cDNA synthesis, primer design and ddPCR .....	8
Progesterone production .....	9
Study Two: Embryo development after COC exposure to 41.0°C for 4 or 6 hIVM.....	10
Statistical analyses .....	10
Chapter Three Results .....	12
Study One: Interleukin 6, IL6 Receptor, Signal Transducer and LIF transcripts during <i>in vitro</i> maturation in COCs matured at 38.5 or 41.0°C .....	12
Relative abundance of IL6 in COCs matured at 38.5 and 41.0°C .....	12
Relative abundance of IL6 receptor (IL6R) in COCs matured at 38.5 and 41.0°C .....	12
Relative abundance of IL6 signal transducer (IL6ST) in COCs matured at 38.5 and 41.0°C .....	12
Relative abundance of leukemia inhibitory factor (LIF) .....	13
Progesterone production .....	13
Study Two: Embryo development after COCs exposed to 41.0°C for 4 or 6 hIVM.....	13
Chapter Four Discussion .....	15
Chapter Five Summary, Conclusions and Research Acknowledgments.....	20
List of References.....	21
Vita .....	39

## CHAPTER ONE

### INTRODUCTION AND REVIEW OF PERTINENT LITERATURE

Dairy cows lose the ability to maintain body temperature when temperature-humidity index approaches 72 [1]. In moderate to severe instances of heat stress, cow body temperature may reach or exceed 41.0°C [2-7]. This level of hyperthermia is problematic as for each 1°C increase in rectal temperature pregnancy rate decreases by ~25% [8]. Hyperthermia occurring at or near the time of breeding is especially problematic. Cows having elevated rectal temperatures before artificial insemination are more likely to return to service and have lower conception rates [9]. Hyperthermia related decreases in fertility are not limited to Holstein cows but are problematic in other breeds [10], including *Bos indicus* cattle [11].

Although mechanisms underlying reduced fertility are multifactorial, and to some extent relate to direct effects of elevated body temperature on the cumulus-oocyte complex (COC; [12-14]); heat-induced alterations in the maternal environment (e.g., ovulatory follicle components) are also problematic. Towards this end, Rispoli et al. [15] examined the follicular fluid proteome of lactating dairy cows that became hyperthermic as a result of an acute heat stress event occurring after a pharmacologically-induced LH surge. Hyperthermic cows had increased intrafollicular levels of IL6 in the ovulatory follicle [15]. Consequences specific to IL6 production were likely limited to ovulatory follicle components because levels in circulation were similar in cows maintained in thermoneutral and heat stress conditions [15]. Interleukin 6 is produced by the mural granulosa cells [murine: 16, porcine: 17] and the cumulus-oocyte complex [murine: 16, bovine: 18, ovine: 19, human: 20, 21].

Although functional significance of hyperthermia-induced increases in intrafollicular IL6 remains unclear, heat-induced increases in other cell types [22-24] along with noted effects to promote meiotic maturation [murine: 16, porcine: 17, bovine: 25] and cumulus expansion [murine: 16, 26] [porcine: 17, human: 27] prompted us to conduct the study described herein to examine *IL6*



abundance in the cumulus-oocyte complex when directly exposed to the physiologically-relevant elevated temperature of 41.0°C throughout *in vitro* maturation. Because functional significance of IL6 is dependent on having receptor components, relative abundance of its receptor and signal transducer was also evaluated. Signal transducer is shared by IL6 and LIF [18, 28] thus LIF transcript abundance was also examined. To determine the extent to which direct exposure of COCs to 41.0°C for the short time periods of 4 or 6 h impacted embryo development, ability of oocytes to cleave and develop to the blastocyst stage after performing *in vitro* fertilization was also examined.

## CHAPTER TWO

### MATERIALS AND METHODS

#### **Collection and *in vitro* maturation of cumulus-oocyte complexes**

Reagents and chemicals were obtained from MilliporeSigma (St. Louis, MO, USA) unless indicated otherwise. Oocytes were collected from abattoir-derived ovaries [13], with modifications noted below. Media were prepared per Rispoli *et al.* [29]. Folltropin-V (FSH) was obtained from VetrepHarm Canada, INC. (London, ON, Canada); same batch was used throughout. Cumulus-oocyte complexes (COCs) with compact cumulus cell vestments and homogenous ooplasm underwent *in vitro* maturation (Study 1: ~30 COCs per 0.5 ml maturation medium in polystyrene tubes; Sarstedt AG and Co., Nümbrecht, Germany; Study 2: 29 to 45 COCs (mean =  $34.3 \pm 0.66$ ) per 0.5 mL in 4-well Nunc culture dishes; Thermo Fisher Scientific, Waltham, MA, USA). Incubator temperatures were verified before and during different studies using mercury thermometers sealed in media-filled bottles.

#### **Study one: Interleukin 6, IL6 Receptor, Signal Transducer and LIF transcripts during *in vitro* maturation in COCs matured at 38.5 or 41.0°C**

Cumulus-oocyte complexes were matured at 38.5 or 41.0°C (exposure to 41.0°C was restricted to first 12 h only; thereafter COCs were transferred to 38.5°C). At 2, 4, 6, 8, 10, 12, 16, 20 and 24 h *in vitro* maturation (hIVM) subsets of COCs were removed from culture and kept separate by treatment (Figure 1; 2 x 9 factorial treatment arrangement). Per each time period, COCs were washed twice in Dulbecco's phosphate buffered saline containing 0.1% polyvinyl alcohol and pelleted (600 x g, 5 min). After supernatant removal, COCs were lysed in extraction buffer (Quick-RNA Kit; Zymo Research, Irvine, CA, USA) and stored at -80°C until RNA isolation. A subset of COCs was also processed soon after removal from ovary to provide a 0 hIVM group. Maturation medium that was conditioned by COCs during culture was centrifuged (5 min, 3000 x g);

supernatant was stored at -20°C. Cumulus-oocyte complexes were collected from ovaries on four different days with 3,840 total COCs being utilized. On a given days collection, two different pools of 30 COCs per each temperature were evaluated at 0, 4, 8, 12, 16, 20, and 24 hIVM resulting in a total of 8 observations per these treatment combinations. Related to 2, 6, and 10 hIVM, only one group of 30 COCs was utilized per these treatment combinations resulting in a total of 4 observations for these time periods.

### ***Total RNA isolation, cDNA synthesis, primer design and ddPCR***

Total COC RNA was isolated using the Quick-RNA Microprep Kit (Zymo Research, Irvine, CA, USA) with on-column DNase treatment per manufacturer. Quantity (Nanodrop ND-1000; NanoDrop Technologies, USA) and quality (RNA Nano LabChip; Bioanalyzer 2100, Agilent, USA) of total RNA were determined (RIN values ranged from 7.2 to 10; median of 8.9). Reverse transcription with oligo (dT) and random primers (500 ng per 20 ul reaction; iScript Reverse Transcription Supermix, Bio-Rad, Hercules, CA, USA) was performed per manufacturer and diluted 5-fold with 1 mM Tris-HCl (pH 8.0) and 0.01 mM EDTA (0.1X TE) before performing digital droplet polymerase chain reactions (ddPCR) analyses. A pool resulting from all samples within each collection day was sham-transcribed (iScript No-RT Control Supermix, Bio-Rad) as an additional control.

Primer-BLAST (National Center for Biotechnology Information; U.S. National Library of Medicine, Bethesda, MD, USA) was utilized to design primers spanning exon-exon junction and/or introns (Table 1). Resulting amplicons were sequenced to ensure specificity. Digital droplet PCR was performed in duplicate using 10 ng of nucleic acid per reaction per manufacturer's instructions. No template controls (NTC; 0.1X TE) were analyzed to assess background signal and control for exogenous contamination. Samples were amplified for 40 cycles, 30s per conditions in Table 1

followed by signal stabilization (4°C for 5 min, 40°C for 5 min, hold at 12°C). Acquired data were analyzed using QuantaSoft Analysis Pro (ver. 1.0, Bio-Rad) to calculate number of copies per  $\mu\text{L}$ .

Transcript abundance was normalized to succinate dehydrogenase A (SDHA). Succinate dehydrogenase A has been used as a normalizer for *in vivo* and *in vitro* matured COCs [30-33] and for heat-stressed COCs [34]. It is stably expressed in cumulus during maturation [30, 35] and not altered by elevated temperature exposure (JL Edwards & J Klabnik, unpublished data).

### ***Progesterone production***

Progesterone released into the maturation medium by COCs matured at 38.5 or 41.0°C (Figure 1) was analyzed by radioimmunoassay per manufacturer's instructions (Double Antibody RIA; MP Biomedicals, Santa Ana, CA., USA). Assay sensitivity was 0.02 ng/mL; inter- and intra-assay coefficients of variation were 7.6 and 6.0%, respectively.

### **Study Two: Embryo development after COC exposure to 41.0°C for 4 or 6 hIVM**

Because heat-induced increases in *IL6* levels were noted by 4 hIVM and by 4 and 6 hIVM for *IL6ST*, a second study was performed to evaluate consequences of a 41.0°C exposure for 4 or 6 h on embryonic development. Cumulus-oocyte complexes meeting criteria described above were randomly allocated to three different treatment groups: 38.5°C for 24 hIVM, 41.0°C for 4 hIVM, or 41.0°C for 6 hIVM. After 4 or 6 hIVM at 41.0°C, COCs were transferred to 38.5°C for remainder of *in vitro* maturation. After a total of 24 hIVM, a combination of frozen-thawed-washed sperm from two bulls was added at ~500,000 motile sperm/ml to each well of COCs. Presumptive zygotes were denuded of cumulus and associated sperm at ~16 to 18 h after addition of sperm. Embryonic cleavage was assessed 66 to 70 h after addition of sperm at which point essential amino acids were added to culture medium. At 172 to 178 h after addition of sperm, blastocyst development was recorded. Blastocyst stage and quality scoring was performed as described by Schrock et al. [36]. Nuclei were enumerated using fluorescent microscopy (40X magnification using a Nikon Eclipse

TE300; UV-2A filter: ex 330 to 380 nm, em 400 to 420 nm; Nikon Instruments, Melville, NY, USA) after fixation in 3% paraformaldehyde. Thereafter, embryos were stained using 5 µg/ml Hoechst 33342, washed, and then mounted on glass slides in Dulbecco's phosphate buffered saline containing 50% glycerol and 0.5 µg/ml Hoechst 33342.

### **Statistical analyses**

For study one, data were analyzed using generalized linear mixed models (PROC GLIMMIX, SAS 9.4, SAS Institute, Cary, NC, USA) as a randomized block design, blocking on replicate (i.e., day of COCs collection). Mindful of the 2 x 9 factorial treatment arrangement, fixed effects in the model included IVM temperature (38.5 and 41.0°C), IVM time (2, 4, 6, 8, 10, 12, 16, 20 and 24 h), and respective interaction (IVM temperature x IVM time; 18 treatment combinations). Treatment differences were determined using F-protected least significant differences and are reported as least squares means  $\pm$  standard error.

Multisource nonlinear mixed model regression (JMP PRO 14, SAS Institute, Cary, NC, USA) was performed *a posteriori* to determine the extent to which maturation of COCs at 41.0°C shifted the timing of changes in transcript abundance compared to levels observed in COCs matured at 38.5°C. The nonlinear prediction model was  $a \cdot \text{Exp}(-(0.5 \cdot ((h\text{IVM}-b)/c)^2))$  where *a* is peak (highest) value, *b* is when peak value occurred, and *c* is growth rate (width of temporal production).

Study two was analyzed as a randomized complete block, blocking on day of oocyte collection with main effect of treatment. Mixed model analysis of variance (PROC GLIMMIX; SAS 9.4) was utilized to determine treatment differences in relation to responses in embryo development including rates of cleavage, 8 to 16 cells, and blastocysts as stage and quality scores. Treatment differences were determined using F-protected least significant differences.

## CHAPTER THREE

### RESULTS

#### **Study One: Interleukin 6, IL6 Receptor, Signal Transducer and LIF transcripts during *in vitro* maturation in COCs matured at 38.5 or 41.0°C**

##### ***Relative abundance of IL6 in COCs matured at 38.5 and 41.0°C***

The differences in *IL6* transcript abundance differed depending upon IVM temperature and hIVM (Temp x hIVM,  $P < 0.0001$ ; Figure 2A). While barely detectable soon after COC collection (0 hIVM), *IL6* abundance at 2 hIVM was similar between COCs matured at 38.5 and 41.0°C. By 4 hIVM, acute exposure to 41.0°C increased relative abundance of *IL6* transcript compared to 38.5°C. Interestingly, by 6 and 8 hIVM the relative abundance of *IL6* in COCs exposed to 41.0°C was lower than that observed in COCs matured at 38.5°C. By 12 hIVM, relative abundance of *IL6* was similar between 41.0 and 38.5°C COCs, levels remained low for remainder of maturation.

Use of multisource nonlinear mixed model regression showed that the major consequence of 41.0°C exposure at the beginning of maturation was to shift the temporal production of *IL6*. To this end, *IL6* levels peaked at 4.67 hIVM in COCs directly exposed to 41.0°C, whereas *IL6* levels peaked at 6.44 hIVM when COCs were matured at 38.5°C ( $P < 0.001$ ; Figure 2A, Table 2). Peak values and growth rate were similar in COCs matured at 38.5 and 41.0°C (Table 2).

##### ***Relative abundance of IL6 receptor (IL6R) in COCs matured at 38.5 and 41.0°C***

Differences in the relative abundance of *IL6R* differed depending on hIVM and IVM temperature (Temp x hIVM,  $P = 0.05$ ; Figure 2B). Abundance of *IL6R* was highest during the first 8 hIVM with abundance decreasing thereafter and reaching lowest levels by 20 to 24 hIVM. Except for the 10 hIVM time period, *IL6R* levels were similar in COCs matured at 38.5 and 41.0°C (Figure 2B, Table 2).

### ***Relative abundance of IL6 signal transducer (IL6ST) in COCs matured at 38.5 and 41.0°C***

Relative abundance of *IL6ST* transcripts differed depending on IVM temperature and hIVM (Temp x hIVM,  $P = 0.002$ ; Figure 2C). While barely detectable in COCs soon after collection from antral follicles, *IL6ST* abundance at 2 hIVM was similar between COCs matured at 38.5 and 41.0°C. However, by 4 and 6 hIVM, 41.0°C exposure resulted in higher levels of *IL6ST* compared to 38.5°C counterparts. At 8 hIVM relative abundance of *IL6ST* in COCs matured at 41.0°C was similar to COCs matured at 38.5°C. By 10 hIVM, *IL6ST* was lower in COCs matured at 41.0°C compared to those matured at 38.5 °C. By 16 hIVM, relative abundance of *IL6ST* was similar between 41.0 and 38.5°C and equivalent to levels observed at the onset of maturation (i.e., 2 hIVM).

Use of multisource nonlinear mixed model regression showed that the major consequence of 41.0°C exposure at the beginning of maturation was to shift the temporal production of *IL6ST*. To this end, *IL6ST* peak levels occurred at 6.95 hIVM when COCs were matured at 41.0°C, whereas peak values were noted at 8.29 hIVM when COCs were matured at 38.5°C ( $P < 0.01$ ; Figure 2C, Table 2). Peak values and growth rates for *IL6ST* were similar in COCs matured at 38.5 and 41.0°C (Table 2).

### ***Relative abundance of leukemia inhibitory factor (LIF)***

Relative abundance of *LIF* transcript was affected by hIVM (time,  $P < 0.0001$ , Figure 3) but not by maturation temperature. Soon after collection and placement of COCs in maturation medium, *LIF* levels increased up through 8 hIVM. After 10 hIVM *LIF* levels decreased. By 12 hIVM, relative abundance of *LIF* was similar to values obtained at 2, 4 and 6 hIVM.

### ***Progesterone production***

Progesterone released per COC into the maturation medium differed depending on IVM temperature and hIVM (Temp x hIVM,  $P = 0.0116$ ; Figure 4A). When all time points were included

in the model (2 to 24 hIVM), heat-induced increases in progesterone produced per COC were most prominent at 20 and 24 hIVM (Figure 4A). When including only the time periods when *IL6* levels were shifted by direct exposure to 41.0°C (i.e., 2 to 8 hIVM), progesterone produced per COC was only affected by IVM temperature and hIVM (i.e., no interaction; heat induced differences were not influenced by time; Figure 4B). To this end, progesterone per COC was 45.7 vs 54.9 pg when COCs were matured at 38.5 and 41.0°C ( $P = 0.002$ ). Independent of temperature but related to time (hIVM), progesterone per COC was 12.7, 40.7, 63.2 and 84.5 pg at 2, 4, 6 and 8 hIVM, respectively ( $P < 0.0001$ ).

#### **Study Two: Embryo development after COCs exposed to 41.0°C for 4 or 6 hIVM**

Ability of COCs to cleave and develop to the blastocyst stage after being exposed to an acute, short-term heat shock of 41.0°C during the first 4 or 6 hIVM was similar to COCs matured at 38.5°C (Table 3). Stage and quality of blastocyst stage embryos from COCs matured at 38.5 or 41.0°C were similar. Number of nuclei in blastocyst stage embryos did not differ when originating from COCs matured at 38.5 °C for 24 h, 41.0 °C for 4hIVM or 41.0°C for 6 hIVM (Table 3).



## CHAPTER FOUR

### DISCUSSION

Novel findings described herein provide further insight related to *IL6* and its receptor signaling component transcripts in the bovine cumulus-oocyte complex as it undergoes maturation. Examination at frequent time intervals confirmed temporal expression of *IL6* and its associated *IL6ST* to the first 12 hours of *in vitro* maturation. Interestingly, the major impact of 41.0°C was to shift the timing of expression for *IL6* and its associated signal transducer. Specific to this end, peak *IL6* levels in COCs exposed to an acute heat shock of 41.0°C occurred ~2 hours *earlier* than thermoneutral COCs. Regarding impact on *IL6ST* expression, peak levels in COCs exposed to 41.0°C occurred ~1.5 hours *earlier* than thermoneutral COCs. Functional significance of these findings along with heat-induced increases in progesterone remain unclear, but heat-related changes may be important to affect/promote meiotic maturation.

Interleukin 6 is a pleiotropic cytokine produced by numerous different cell types [37-39]. Specific to the COC, both the oocyte and its associated cumulus produce IL6 (bovine: [18], human: [20, 21], murine: [16], ovine: [19]. Our study demonstrated that the *IL6* transcript is minimally detectable in germinal vesicle (GV) stage COCs soon after removal from antral follicles. Transcript levels are higher by 2 hIVM and peak 6.44 hours in COCs matured 38.5°C; levels decrease thereafter and remain low for remainder of maturation. Collectively these findings document temporal expression of *IL6* in the bovine during the first part of maturation and extend the findings of Tscherner et al. [18] who examined *IL6* abundance in bovine COCs at 0, 7, and 24 hIVM. In that study [18] *IL6* was abundant at 7 hIVM but was not apparent in COCs soon after removal from antral follicles (0 hIVM) or after 24 hIVM. Temporal expression of IL6 during the first part of maturation occurs in other species. Consistent with our findings in the bovine, *IL6* was detectable but present in low levels in ovine COCs soon after removal from antral follicles [19]. Peak *IL6* was noted at 4 hIVM; levels decreased thereafter. Use of Western blot confirmed presence of the IL6

protein in ovine COCs soon after removal from the antral follicle with highest levels at 4 and 8 hIVM [19].

Regarding the potential for COC-derived *IL6* to be impactful at the level of the cumulus-oocyte complex, outcomes of study one show that *IL6* receptor abundance is greatest when *IL6* levels are markedly increasing during the first 6 hIVM. Furthermore, receptor levels are lowest during the latter half of maturation (~12 h to 24 hIVM) when *IL6* transcripts remain unchanged and are at low levels. Transcript abundance for the *IL6* receptor relates well to protein levels in ovine COCs [19] which has been localized to the oocyte surface [19]. Although receptor binding is important, intracellular signaling related to IL6 is dependent on forming a complex with IL6 signal transducer [40-43]. Like *IL6*, *IL6ST* was temporally expressed during the first half of maturation, except peak levels occurred ~2 hours later. In murine COCs and hybridoma cells, addition of IL6 increases IL6ST transcript and protein levels [16, 44]. Whether or not COC-derived increases in *IL6* affect *IL6ST* expression in the bovine is unclear. Because of its importance after receptor binding, it is intuitive for *IL6ST* levels to peak and persist a bit longer than *IL6*, which is what was observed in study one.

Nonetheless, the presence of this pleiotropic cytokine, its receptor, and associated signal transducer set the stage for IL6 to play an active role within the maturing COC. Towards this end, Liu [16] showed that the addition of IL6 to murine COCs during in vitro maturation improved success of embryo transfers by increasing number of pups born [16]. Other efforts using ovine [19], bovine [abstract: 25] and porcine [abstract:17] COCs noted improvements in meiotic progression (i.e., metaphase I or metaphase II) with the IL6 addition to the maturation medium. “Pro”-effects of IL6 are likely “dose” dependent. Zhao et al. [19] showed that a lower dose of IL6 (10 ng/mL) increased maturation rates, whereas a higher dose (100 ng/mL) reduced maturation rates and impaired embryo development when IL6 was added to ovine COCs during *in vitro* maturation.

Although the specific factor(s) underlying the beginnings of maturation are not yet fully elucidated, an increase in COC-derived *IL6* with levels peaking at or around 6 hIVM is likely a significant promotant of GVBD. In other cell types, IL6 reduces gap junction permeability [45] which is requisite for GVBD. When murine COCs are cultured in a hypoxanthine-containing medium to inhibit spontaneous breakdown of the GV, addition of IL6 and its soluble receptor *induced* GVBD, despite COCs being held in the continued presence of a meiotic inhibitor [16]. Marked increases in COC-derived *IL6* peaking at or around 6 hIVM in study one overlap with time period leading up to and when GVBD occurs [46-49]. Interestingly and consistent with consequences of adding IL6 to meiotically inhibited oocytes, direct exposure of COCs to an elevated temperature of 41.0°C increases IL6 in other cell types [22-24] and *induces* GVBD in bovine COCs [46, 47, 49]. Heat-induced hastening of this developmentally-critical event is detectable as early as 4 hIVM and more prominent by 6 hIVM (Hooper et al., 2015). Recent efforts of Campen et al. [46] demonstrated changes in maternal chromatin at the beginnings of GVBD coincided with heat-induced reductions in gap junction permeability. Although factors triggering accelerated GVBD when activated by 41.0°C remain unclear, novel findings in this study clearly indicating a heat-induced *shift* in the timing of *IL6* expression and its associated signal transducer by ~1.5 to 2 hours *earlier*, attests to the notion for IL6 to be an underlying contributive factor in the heat-induced hastening of GVBD previously reported by others [46, 47, 49].

Leukemia inhibitory factor is a member of the IL6 family [50-53]. Because it is known to positively impact oocyte maturation in multiple species [54-56], examination of transcript abundance throughout maturation complimented efforts described herein examining COC levels of *IL6*. Unlike *IL6*, however, there was no impact of 41.0°C on *LIF* expression at any time period examined. Although *LIF* levels increased during the first part of maturation (up through 8 to 10 hIVM) and decreased thereafter, highest levels were reached ~2 hours *after* *IL6* peaked which is more consistent with changes in *IL6ST*. Interleukin 6 and LIF share the same signal transducer

with IL6 having higher affinity when both ligands are present [18, 28]. Depending on the importance of IL6 as a GVBD promotant (peaks earlier than IL6ST), temporal production of *LIF* to better coincide with signal transducer may be important to influence other developmentally important events for maturation success (e.g., metaphase I and metaphase II progression).

Associated cumulus cells revert back to granulosa cell-like state while the oocyte undergoes maturation [57]. Towards this end and within a short order of time, cumulus-derived progesterone released into the maturation medium increases and continues to do so throughout maturation [46, study one: 58]. When examining just the time periods when *IL6* levels were shifted by direct exposure to 41.0°C (i.e., 2 to 8 hIVM), COCs released more progesterone into maturation medium (45.7 vs 54.9 pg for control and heat stress, respectively). Similar findings were previously reported by Campen et al. [46]. Mindful that levels are likely much higher in the COC, the presence of genomic and non-genomic progesterone receptors in bovine oocytes [59] highlights opportunities for progesterone to be acting in a paracrine manner at the level of the oocyte. Although significance of heat-induced “heightening” of progesterone in the COC remains to be determined, progesterone stimulates GVBD [60] and affects meiotic progression [61] attesting to functional significance during this early time period. In porcine COCs, suppression of progesterone production with aminoglutethimide almost entirely prevented GV-breakdown; inhibitory effects on GV-breakdown were overcome by additional progesterone [62]. Regarding the potential interplay of progesterone and IL6, blocking progesterone’s ability to bind to its receptor using RU486 prevented FSH-induction of *IL6* in murine oocytes [16] suggesting that progesterone may be a contributory factor helping modulate IL6 production.

Exposure to 41.0°C at the onset of maturation may be “shocking” but is unavoidable when striving to understand consequences of elevated temperature(s) when occurring during oocyte maturation. Collection of COCs during late fall, winter, spring, first part of summer before summer heat stress attests to the relevance of our model for gaining an understanding of the consequences

of elevated body temperature(s) that may occur in sexually active females. Depending on management, it is not uncommon for the body temperature of sexually active females to become elevated (up to 1.3°C; [63-70]) with peak levels occurring at or around the time of the LH surge [70-72]. In two different studies described in a manuscript that is in the final stages of preparation for peer review (Edwards et al., unpublished), rectal temperatures of *Bos taurus* (n = 427) and *Bos indicus* (n = 1,278) females submitted for fixed timed artificial insemination ranged from 37.8 to 41.8°C and 37.0 to 40.9°C, respectively. Use of logistic regression showed that per each °C increase in rectal temperature at insemination odds for pregnancy increased by 1.6 and 1.9 times (Edwards et al., unpublished). Mindful of this, but also expecting that prolonged exposures are detrimental to embryo development [12-14, 36, 49, 73-78], additional effort was put forth to examine consequences of 41.0°C exposure when occurring during the first 4 or 6 hIVM. Study two outcomes clearly illustrate that blastocyst development after performing IVF was not affected by 41.0°C exposure for 4 or 6 h. In fact, blastocyst development, stage and quality scores, and nuclei numbers were numerically higher when COCs were matured at 41.0°C versus thermoneutral temperature of 38.5°C. Because the major consequence of 41.0°C was to shift temporal production of *IL6* (study one) and hasten onset of GVBD by an equivalent amount of time, it is unclear if performing IVF earlier than 24 hIVM would have been beneficial as it is when 41.0°C exposure persists for 12 h [36].

## **CHAPTER FIVE**

### **RESEARCH ACKNOWLEDGMENTS**

This project was supported by Agriculture and Food Research Initiative Competitive Grant no. 2016-67015-24899 from the USDA National Institute of Food and Agriculture, the state of Tennessee through UT AgResearch, East Tennessee Research and Education Center, the Department of Animal Science, and the USDA National Institute of Food and Agriculture, Hatch Project No. 1015707 and 1022068. The authors would like to thank Xiaocun Sun from Research Computing Support in OIT at the University of Tennessee for her assistance with JMP analysis. The authors would like to thank Hunter Liles, Abigayle Pollock, and Mary Ali Oliver for their time spent in travel to the abattoir and for collections. The authors also wish to thank DeSoto Biosciences for providing ovaries to obtain COCs for primer optimization.

## **LIST OF REFERENCES**

1. **Armstrong DV.** Heat stress interaction with shade and cooling. *Journal of Dairy Science* 1994; 77: 2044-2050.
2. **Gaalaas R.** Effect of atmospheric temperature on body temperature and respiration rate of Jersey cattle. *Journal of Dairy Science* 1945; 28: 555-563.
3. **Seath DM, Miller GD.** The Relative Importance of High Temperature and High Humidity as Factors Influencing Respiration Rate, Body Temperature, and Pulse Rate of Dairy Cows. *Journal of Dairy Science* 1946; 29: 465-472.
4. **Roman-Ponce H, Thatcher W, Buffington D, Wilcox C, Van Horn H.** Physiological and production responses of dairy cattle to a shade structure in a subtropical environment. *Journal of Dairy Science* 1977; 60: 424-430.
5. **Turner H.** Genetic variation of rectal temperature in cows and its relationship to fertility. *Animal Science* 1982; 35: 401-412.
6. **Elvinger F, Hansen P, Natzke R.** Modulation of function of bovine polymorphonuclear leukocytes and lymphocytes by high temperature in vitro and in vivo. *American Journal of Veterinary Research* 1991; 52: 1692.
7. **Ealy AD, Drost M, Hansen PJ.** Developmental changes in embryonic resistance to adverse effects of maternal heat stress in cows. *Journal of Dairy Science* 1993; 76: 2899-2905.
8. **Ulberg LC, Burfening PJ.** Embryo death resulting from adverse environment on spermatozoa or ova. *Journal of Animal Science* 1967; 26: 571-577.
9. **Fallon G.** Body temperature and fertilization in the cow. *Reproduction* 1962; 3: 116-123.
10. **Dunlap S, Vincent C.** Influence of postbreeding thermal stress on conception rate in beef cattle. *Journal of Animal Science* 1971; 32: 1216-1218.
11. **Zakari A, Molokwu E, Osori D.** Effects of rectal and ambient temperatures and humidity on conception rates. *Theriogenology* 1981; 16: 331-336.



12. **Edwards JL, Hansen P.** Elevated temperature increases heat shock protein 70 synthesis in bovine two-cell embryos and compromises function of maturing oocytes. *Biology of Reproduction* 1996; 55: 340-346.
13. **Lawrence JL, Payton RR, Godkin JD, Saxton AM, Schrick FN, Edwards JL.** Retinol improves development of bovine oocytes compromised by heat stress during maturation. *Journal of Dairy Science* 2004; 87: 2449-2454.
14. **Zhandi M, Towhidi A, Nasr-Esfahani MH, Eftekhari-Yazdi P, Zare-Shahneh A.** Unexpected detrimental effect of insulin like growth factor-1 on bovine oocyte developmental competence under heat stress. *Journal of Assisted Reproduction and Genetics* 2009; 26: 605-611.
15. **Rispoli LA, Edwards JL, Pohler KG, Russell S, Somiari RI, Payton RR, Schrick FN.** Heat-induced hyperthermia impacts the follicular fluid proteome of the periovulatory follicle in lactating dairy cows. *PLOS ONE* 2019; 14.
16. **Liu Z, de Matos DG, Fan H-Y, Shimada M, Palmer S, Richards JS.** Interleukin-6: an autocrine regulator of the mouse cumulus cell-oocyte complex expansion process. *Endocrinology* 2009; 150: 3360-3368.
17. **Faundez R, Kawecka O, Aniolek O, Petrajtis-Golobów M, Karska D, Gajewski Z.** The influence of GM-CSF, IL-6 and TGF- $\alpha$  on in vitro maturation of porcine oocytes: P65. *Reproduction in Domestic Animals* 2015; 50: 51-52.
18. **Tscherner A, Brown AC, Stalker L, Kao J, Dufort I, Sirard M-A, LaMarre J.** STAT3 signaling stimulates miR-21 expression in bovine cumulus cells during in vitro oocyte maturation. *Scientific Reports* 2018; 8: 11527.
19. **Zhao X-a, Cang M, Gao X-y, Yang M-l, Yuan J-l, Zhu B, Wang Z-g, Liu D-j.** Expression of interleukin-6 and interleukin-6 receptor in ovine oocytes during in vitro maturation. *Journal of Integrative Agriculture* 2012; 11: 1333-1339.

20. **Zolti M, Ben-Rafael Z, Meirom R, Shemesh M, Bider D, Mashiach S, Apte RN.**  
Cytokine involvement in oocytes and early embryos. *Fertility and Sterility* 1991; 56: 265-272.
21. **Machelon V, Emilie D, Lefevre A, Nome F, Durand-Gasselin I, Testart J.**  
Interleukin-6 biosynthesis in human preovulatory follicles: some of its potential roles at ovulation. *The Journal of Clinical Endocrinology & Metabolism* 1994; 79: 633-642.
22. **Bharati J, Dangi SS, Mishra SR, Chouhan VS, Verma V, Shankar O, Bharti MK, Paul A, Mahato DK, Rajesh G, Singh G, Maurya VP, Bag S, Kumar P, Sarkar M.**  
Expression analysis of toll like receptors and interleukins in tharparkar cattle during acclimation to heat stress exposure. *Journal of Thermal Biology* 2017; 65: 48-56.
23. **Obi S, Nakajima T, Hasegawa T, Kikuchi H, Oguri G, Takahashi M, Nakamura F, Yamasoba T, Sakuma M, Toyoda S, Tei C, Inoue T.** Heat induces interleukin-6 in skeletal muscle cells via TRPV1/PKC/CREB pathways. *Journal of Applied Physiology* 2016; 122: 683-694.
24. **Welc SS, Phillips NA, Oca-Cossio J, Wallet SM, Chen DL, Clanton TL.**  
Hyperthermia increases interleukin-6 in mouse skeletal muscle. *American Journal of Physiology-Cell Physiology* 2012; 303: C455-C466.
25. **Faundez R, Kawecka O, Karska D, Petrajtis M, Gajewski Z.** The influence of GM-CSF, IL-6 AND TGF- $\alpha$  on the in vitromaturation of bovine oocytes: P63. *Reproduction in Domestic Animals* 2014; 49.
26. **Wang Y, Liang N, Yao G, Tian H, Zhai Y, Yin Y, Sun F.** Knockdown of TrkA in cumulus oocyte complexes (COCs) inhibits EGF-induced cumulus expansion by down-regulation of IL-6. *Molecular and Cellular Endocrinology* 2014; 382: 804-813.
27. **Clark A.** Use of IL-6 type cytokines for maturation of oocytes. United States: Merek Serono SA; 2011.

28. **Gearing D, Thut C, VandeBos T, Gimpel S, Delaney P, King J, Price V, Cosman D, Beckmann M.** Leukemia inhibitory factor receptor is structurally related to the IL-6 signal transducer, gp130. *The EMBO journal* 1991; 10: 2839-2848.
29. **Rispoli LA, Lawrence JL, Payton RR, Saxton AM, Schrock GE, Schrick FN, Middlebrooks BW, Dunlap JR, Parrish JJ, Edwards JL.** Disparate consequences of heat stress exposure during meiotic maturation: embryo development after chemical activation vs fertilization of bovine oocytes. *Reproduction* 2011; 142: 831-843.
30. **Mourad A, Steph JD, Marc-André S.** Cumulus cell gene expression following the LH surge in bovine preovulatory follicles: potential early markers of oocyte competence. *Reproduction* 2010; 140: 835-852.
31. **del Collado M, da Silveira JC, Sangalli JR, Andrade GM, Sousa LRdS, Silva LA, Meirelles FV, Perecin F.** Fatty acid binding protein 3 and transzonal projections are involved in lipid accumulation during in vitro maturation of bovine oocytes. *Scientific Reports* 2017; 7: 2645.
32. **Botigelli RC, Razza EM, Pioltine EM, Fontes PK, Schwarz KRL, Leal CLV, Nogueira MFG.** Supplementing in vitro embryo production media by NPPC and sildenafil affect the cytoplasmic lipid content and gene expression of bovine cumulus-oocyte complexes and embryos. *Reproductive Biology* 2018; 18: 66-75.
33. **Macabelli CH, Ferreira RM, Gimenes LU, de Carvalho NAT, Soares JG, Ayres H, Ferraz ML, Watanabe YF, Watanabe OY, Sangalli JR, Smith LC, Baruselli PS, Meirelles FV, Chiaratti MR.** Reference gene selection for gene expression analysis of oocytes collected from dairy cattle and buffaloes during winter and summer. *PLOS ONE* 2014; 9: e93287.

34. **Pavani KC, Rocha A, Baron E, Lourenço J, Faheem M, da Silva FM.** The effect of kinetic heat shock on bovine oocyte maturation and subsequent gene expression of targeted genes. *Zygote* 2017; 25: 383-389.
35. **Regassa A, Rings F, Hoelker M, Cinar U, Tholen E, Looft C, Schellander K, Tesfaye D.** Transcriptome dynamics and molecular cross-talk between bovine oocyte and its companion cumulus cells. *BMC Genomics* 2011; 12: 57.
36. **Schrock G, Saxton A, Schrick F, Edwards J.** Early in vitro fertilization improves development of bovine ova heat stressed during in vitro maturation. *Journal of Dairy Science* 2007; 90: 4297-4303.
37. **Brown HM, Dunning KR, Sutton-McDowall M, Gilchrist RB, Thompson JG, Russell DL.** Failure to launch: aberrant cumulus gene expression during oocyte in vitro maturation. *Reproduction (Cambridge, England)* 2016; 153: R109-R120.
38. **Schett G.** Physiological effects of modulating the interleukin-6 axis. *Rheumatology* 2018; 57: ii43-ii50.
39. **Smolikova K, Mlynarcikova A, Scsukova S.** Role of interleukins in the regulation of ovarian functions. *Endocrine Regulations* 2012; 46: 237-253.
40. **HEINRICH PC, BEHRMANN I, HAAN S, HERMANN S, MÜLLER-NEWEN G, SCHAPER F.** Principles of interleukin (IL)-6-type cytokine signalling and its regulation. *Biochemical Journal* 2003; 374: 1-20.
41. **Hibi M, Murakami M, Saito M, Hirano T, Taga T, Kishimoto T.** Molecular cloning and expression of an IL-6 signal transducer, gp130. *Cell* 1990; 63: 1149-1157.
42. **Wolf J, Rose-John S, Garbers C.** Interleukin-6 and its receptors: a highly regulated and dynamic system. *Cytokine* 2014; 70: 11-20.

43. **Mackiewicz A, Schooltink H, Heinrich P, Rose-John S.** Complex of soluble human IL-6-receptor/IL-6 up-regulates expression of acute-phase proteins. *The Journal of Immunology* 1992; 149: 2021-2027.
44. **Canellada A, Alvarez I, Berod L, Gentile T.** Estrogen and progesterone regulate the IL-6 signal transduction pathway in antibody secreting cells. *The Journal of Steroid Biochemistry and Molecular Biology* 2008; 111: 255-261.
45. **Temme A, Traub O, Willecke K.** Downregulation of connexin32 protein and gap-junctional intercellular communication by cytokine-mediated acute-phase response in immortalized mouse hepatocytes. *Cell and Tissue Research* 1998; 294: 345-350.
46. **Campen KA, Abbott CR, Rispoli LA, Payton RR, Saxton AM, Edwards JL.** Heat stress impairs gap junction communication and cumulus function of bovine oocytes. *Journal of Reproduction and Development* 2018; 64: 385 - 392.
47. **Hooper LM, Payton RR, Rispoli LA, Saxton AM, Edwards JL.** Impact of heat stress on germinal vesicle breakdown and lipolytic changes during in vitro maturation of bovine oocytes. *The Journal of Reproduction and Development* 2015; 61: 459-464.
48. **Hyttel P, Xu K, Smith S, Greve T.** Ultrastructure of in-vitro oocyte maturation in cattle. *Reproduction* 1986; 78: 615-625.
49. **Edwards J, Saxton A, Lawrence J, Payton R, Dunlap J.** Exposure to a physiologically relevant elevated temperature hastens in vitro maturation in bovine oocytes. *Journal of dairy science* 2005; 88: 4326-4333.
50. **Turnley AM, Bartlett PF.** Cytokines that signal through the leukemia inhibitory factor receptor- $\beta$  complex in the nervous system. *Journal of Neurochemistry* 2000; 74: 889-899.
51. **Fourcin M, Chevalier S, Lebrun JJ, Kelly P, Pouplard A, Wijdenes J, Gascan H.** Involvement of gp130/interleukin-6 receptor transducing component in interleukin-11 receptor. *European Journal of Immunology* 1994; 24: 277-280.

52. **Kishimoto T, Taga T, Akira S.** Cytokine signal transduction. *Cell* 1994; 76: 253-262.
53. **Ip NY, Nye SH, Boulton TG, Davis S, Taga T, Li Y, Birren SJ, Yasukawa K, Kishimoto T, Anderson DJ.** CNTF and LIF act on neuronal cells via shared signaling pathways that involve the IL-6 signal transducing receptor component gp130. *Cell* 1992; 69: 1121-1132.
54. **Dang-Nguyen TQ, Haraguchi S, Kikuchi K, Somfai T, Bodó S, Nagai T.** Leukemia inhibitory factor promotes porcine oocyte maturation and is accompanied by activation of signal transducer and activator of transcription 3. *Molecular Reproduction and Development* 2014; 81: 230-239.
55. **Mo X, Wu G, Yuan D, Jia B, Liu C, Zhu S, Hou Y.** Leukemia inhibitory factor enhances bovine oocyte maturation and early embryo development. *Molecular Reproduction and Development* 2014; 81: 608-618.
56. **Wang J, Liu Z, Sun Q, Xia S, Cui J, Yang L, An L, Zhang J, Su L, Su Y, Du F.** Combined treatment with cysteamine and leukemia inhibitory factor promotes guinea pig oocyte meiosis in vitro. *American Journal of Translational Research* 2019; 11: 7479-7491.
57. **Chaffin CL, Lee YS, VandeVoort CA, Patel BG, Latham KE.** Rhesus monkey cumulus cells revert to a mural granulosa cell state after an ovulatory stimulus. *Endocrinology* 2012; 153: 5535-5545.
58. **Rispoli LA, Payton RR, Gondro C, Saxton AM, Nagle KA, Jenkins BW, Schrick FN, Edwards JL.** Heat stress effects on the cumulus cells surrounding the bovine oocyte during maturation: altered matrix metalloproteinase 9 and progesterone production. *Reproduction* 2013; 146: 193-207.

59. **Aparicio I, Garcia-Herreros M, O'shea L, Hensey C, Lonergan P, Fair T.** Expression, regulation, and function of progesterone receptors in bovine cumulus oocyte complexes during in vitro maturation. *Biology of Reproduction* 2011; 84: 910-921.
60. **Siqueira LC, Barreta MH, Gasperin B, Bohrer R, Santos JT, Junior JB, Oliveira JF, Gonçalves PB.** Angiotensin II, progesterone, and prostaglandins are sequential steps in the pathway to bovine oocyte nuclear maturation. *Theriogenology* 2012; 77: 1779-1787.
61. **Sirotkin AV.** Involvement of steroid hormones in bovine oocytes maturation in vitro. *The Journal of Steroid Biochemistry and Molecular Biology* 1992; 41: 855-858.
62. **Shimada M, Terada T.** FSH and LH induce progesterone production and progesterone receptor synthesis in cumulus cells: a requirement for meiotic resumption in porcine oocytes. *Molecular Human Reproduction* 2002; 8: 612-618.
63. **Piccione G, Caola G, Refinetti R.** Daily and estrous rhythmicity of body temperature in domestic cattle. *BMC Physiology* 2003; 3: 1-8.
64. **Suthar V, Burfeind O, Patel J, Dhami A, Heuwieser W.** Body temperature around induced estrus in dairy cows. *Journal of Dairy Science* 2011; 94: 2368-2373.
65. **Higaki S, Miura R, Suda T, Andersson LM, Okada H, Zhang Y, Itoh T, Miwakeichi F, Yoshioka K.** Estrous detection by continuous measurements of vaginal temperature and conductivity with supervised machine learning in cattle. *Theriogenology* 2019; 123: 90-99.
66. **Randi F, McDonald M, Duffy P, Kelly AK, Lonergan P.** The relationship between external auditory canal temperature and onset of estrus and ovulation in beef heifers. *Theriogenology* 2018; 110: 175-181.

67. **Miura R, Yoshioka K, Miyamoto T, Nogami H, Okada H, Itoh T.** Estrous detection by monitoring ventral tail base surface temperature using a wearable wireless sensor in cattle. *Animal Reproduction Science* 2017; 180: 50-57.
68. **Kyle B, Kennedy A, Small J.** Measurement of vaginal temperature by radiotelemetry for the prediction of estrus in beef cows. *Theriogenology* 1998; 49: 1437-1449.
69. **Lewis G, Newman S.** Changes throughout estrous cycles of variables that might indicate estrus in dairy cows. *Journal of Dairy Science* 1984; 67: 146-152.
70. **Fisher A, Morton R, Dempsey J, Henshall J, Hill J.** Evaluation of a new approach for the estimation of the time of the LH surge in dairy cows using vaginal temperature and electrodeless conductivity measurements. *Theriogenology* 2008; 70: 1065-1074.
71. **Rajamahendran R, Robinson J, Desbottes S, Walton JS.** Temporal relationships among estrus, body temperature, milk yield, progesterone and luteinizing hormone levels, and ovulation in dairy cows. *Theriogenology* 1989; 31: 1173-1182.
72. **Mosher M, Ottobre J, Haibel G, Zartman D.** Estrual rise in body temperature in the bovine II. The temporal relationship with ovulation. *Animal Reproduction Science* 1990; 23: 99-107.
73. **Roth Z, Hansen P.** Sphingosine 1-phosphate protects bovine oocytes from heat shock during maturation. *Biology of Reproduction* 2004; 71: 2072-2078.
74. **Roth Z, Hansen P.** Involvement of apoptosis in disruption of developmental competence of bovine oocytes by heat shock during maturation. *Biology of Reproduction* 2004; 71: 1898-1906.
75. **de Castro e Paula LA, Hansen P.** Interactions between oxygen tension and glucose concentration that modulate actions of heat shock on bovine oocytes during in vitro maturation. *Theriogenology* 2007; 68: 763-770.



76. **Sugiyama S, McGowan M, Phillips N, Kafi M, Young M.** Effects of increased ambient temperature during IVM and/or IVF on the in vitro development of bovine zygotes. *Reproduction in Domestic Animals* 2007; 42: 271-274.
77. **Edwards J, Bogart A, Rispoli L, Saxton A, Schrick F.** Developmental competence of bovine embryos from heat-stressed ova. *Journal of Dairy Science* 2009; 92: 563-570.
78. **Soto P, Smith LC.** BH4 peptide derived from Bcl-xL and Bax-inhibitor peptide suppresses apoptotic mitochondrial changes in heat stressed bovine oocytes. *Molecular Reproduction and Development* 2009; 76: 637-646.

Table 1: Primer sequences and annealing conditions used for ddPCR

Gene	GenBank Accession Number	Amplicon Location (bp)	Primer Set	Primer Concentration (nM)	Annealing Temperature (°C)
<i>IL6</i>	NM_173923.2	349-568	3'-GCATCTTCTCCAGCAGGTCAG 5'-CAATCTGGGTTCAATCAGGCGAT	250	56
<i>IL6R</i>	NM_001110785.3	343-666	3'-TCGGGCTGTAGGAGTTTGTAGC 5'-GCGCTTGGTGGTGGATGTTC	125	56
<i>IL6ST</i>	XM_010816769.3*	1136-1355	3'-CGCGTCTGATTTGCCAACAA 5'-GTCTCATGCTCACGGCACTA	250	58
<i>LIF</i>	NM_173931	157-359	3'-CTGGGCCGTGTAATAGAGGAT 5'-TCTTGGCGGCAGGAGTTGT	250	58
<i>SDHA</i>	NM_174178	1433-1646	3'-TCCGTAGAGGCTGCTGATCT 5'-GTCCTGCAGACCCGGAGATA	250	58

\* Wooldridge et al. (2019)

Table 2: Impact of 41.0°C exposure on IL6, IL6R and IL6ST levels in COCs during *in vitro* maturation using multisource nonlinear mixed model regression.

Transcript		Peak Value*	Peak Time (hIVM)	Growth Rate**
<i>IL6</i>	38.5°C	0.32 ± 0.03 <sup>a</sup>	6.44 ± 0.23 <sup>a</sup>	2.43 ± 0.28 <sup>a</sup>
	41.0°C	0.33 ± 0.03 <sup>a</sup>	4.67 ± 0.23 <sup>b</sup>	1.82 ± 0.22 <sup>a</sup>
	P-value	P > 0.05	<b>P &lt; 0.001</b>	P > 0.05
	R <sup>2</sup>	0.67		
	SSE	0.41		
<i>IL6R</i>	38.5°C	0.01 ± 0.00 <sup>a</sup>	5.02 ± 1.49 <sup>a</sup>	6.33 ± 2.01 <sup>a</sup>
	41.0°C	0.01 ± 0.00 <sup>a</sup>	5.16 ± 0.70 <sup>a</sup>	4.42 ± 0.82 <sup>a</sup>
	P-value	P > 0.05	P > 0.05	P > 0.05
	R <sup>2</sup>	0.33		
	SSE	0.001		
<i>IL6ST</i>	38.5°C	2.66 ± 0.19 <sup>a</sup>	8.29 ± 0.23 <sup>a</sup>	2.66 ± 0.22 <sup>a</sup>
	41.0°C	2.77 ± 0.20 <sup>a</sup>	6.95 ± 0.23 <sup>b</sup>	2.63 ± 0.23 <sup>a</sup>
	P-value	P > 0.05	<b>P &lt; 0.01</b>	P > 0.05
	R <sup>2</sup>	0.69		
	SSE	22.7		

\*Peak value: highest obtained level

\*\*Growth rate: Full Width Half Maximum (3 standard deviations from the mid-point at half maximum)

<sup>a,b</sup> means differ P < 0.05

Table 3: Impact of an acute exposure to 41.0°C for the first 4 or 6 h of *in vitro* maturation

Treatment	No. OMM <sup>1</sup>	Cleaved (%)	8 to 16-cell (%)	Blastocysts (%)	Stage	Quality	Nuclei
38.5°C-24 h	454	68.87 ± 3.48	74.64 ± 3.47	23.81 ± 2.77	6.69 ± 0.12	1.79 ± 0.12	101.68 ± 11.39
41.0°C-4 h*	444	74.77 ± 3.19	75.19 ± 3.39	29.61 ± 3.08	6.75 ± 0.12	1.96 ± 0.12	118.76 ± 11.39
41.0°C-6 h*	440	75.62 ± 3.07	65.88 ± 3.78	27.73 ± 2.92	6.64 ± 0.12	1.74 ± 0.12	116.07 ± 11.39
	<i>P-value</i>	0.1426	0.0714	0.2558	0.6006	0.1586	0.4313

\*Exposed to 41.0°C for 4 or 6 h at onset of maturation period followed by 38.5°C for a total of 24 h

<sup>1</sup>Number of COCs placed in maturation medium (OMM) for indicated treatment




Figure 1: Schematic of study design. At 2, 4, 6, 8, 10, 12, 16, 20, or 24 hIVM subsets of COCs matured at 38.5°C (designated by white circles) or 41.0°C (first 12 designated by black circles, then moved to 38.5°C designated by gray circles) were removed from culture, washed, lysed before storage at -80°C until RNA extraction and subsequent RNA analyses.




Figure 2: Relative abundance of interleukin 6 and signaling molecules in cumulus-oocyte complexes matured for up to 24 h at either 38.5°C or 41.0°C (first 12 h; 38.5°C thereafter). Interaction of temperature x hIVM between 38.5°C and 41.0°C for *IL6* (Panel A), *IL6R* (Panel B), *IL6ST* (Panel C). Bars (least squares means  $\pm$  SEM) having <sup>A-1</sup> different letter designations differ at indicated *P* (Temp x hIVM). Dashed (38.5°C) and solid (41.0°C) lines indicate relative abundance curves over the first 12 hIVM.




Figure 3: Relative abundance of *LIF* in cumulus-oocyte complexes during maturation, 0 hIVM not included in analysis but as a visual representation of a starting point, averaged across maturation temperatures presented as least squares means  $\pm$  SEM.

<sup>A-D</sup> means differ  $P < 0.05$




Figure 4: Average progesterone produced per cumulus-oocyte complex (COC) during *in vitro* maturation (IVM) at 38.5°C or 41.0°C as measured in conditioned medium. COCs underwent IVM for up to 24 h at 38.5°C or 41.0°C (first 12 hours; 38.5°C thereafter). (A) Temperature x hIVM P = 0.012; \* denotes heat-induced increase in progesterone at indicated time point (B) Impact of 41.0°C on COC on progesterone production during time period when relative abundance of *IL6* was altered by 41.0°C.



## **VITA**

Julia Rose Rowinski was born in Lemont, Illinois. She attended University of Kentucky where she received her B.S. in Animal Science with a minor in Biology. She was a member of Sigma Alpha, block and bridle, and UK shotgun team at University of Kentucky. Julia was raised going to work with her mother at an human infertility clinic and that is where her interest for IVF and reproduction stemmed from. In the fall of 2018, she arrived at the University of Tennessee to pursue her M.S. in reproductive physiology under the mentorship of Dr. Edwards.